



Development of a video-microscopic method to compare the effect of precipitation inhibitors

Christfort, J. F. ; Plum, J.; Madsen, C. M. ; Nielsen, Line Hagner; Müllertz, A.; Rades, T.

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Christfort, J. F., Plum, J., Madsen, C. M., Nielsen, L. H., Müllertz, A., & Rades, T. (2016). *Development of a video-microscopic method to compare the effect of precipitation inhibitors*. Abstract from 10th Annual Meeting of the Pharmaceutical Solid State Research Cluster, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Development of a video-microscopic method to compare the effect of precipitation inhibitors

J. F. Christfort¹, J. Plum¹, C. M. Madsen¹, L. H. Nielsen², A. Müllertz^{1,3}, and T. Rades¹

¹Department of Pharmacy, University of Copenhagen, Universitetsparken 2, 2100 Copenhagen Ø, Denmark

²Department of Micro- and Nanotechnology, Technical University of Denmark, Ørstedes Plads, 2800 Kgs. Lyngby, Denmark

³Bioneer:FARMA, University of Copenhagen, Universitetsparken 2, 2100 Copenhagen Ø, Denmark

PURPOSE

The aim of this study was to develop a microscopic method to evaluate the effect of precipitation inhibitors (PI) on supersaturated solutions of poorly soluble drugs using a digital microscopy imaging system.

METHODS

A supersaturated tadalafil solution was induced by spiking 30 µL of a 5 mg/mL DMSO stock into 200 µL fasted state simulated intestinal medium (FaSSIF, biorelevant.com) with predissolved HPMC (0-1 %w/v) as PI. The studies were performed on the oCelloScope SystemTM (Philips Biocell A/S, Allerød, DK) in a 96-well plate using video-microscopy and the precipitation was followed over time. The induction time was determined as the time point with the first particle in focus and the onwards growth of the particles was measured.

RESULTS

Without PI, the induction time was found to be 5.5 ± 1.1 minutes for tadalafil in FaSSIF. An increase in the induction time was demonstrated with increasing concentrations of HPMC, and with concentrations above 0.05 %w/v, a significantly prolonged induction time was observed (9.0 ± 1.7 to 13.6 ± 1.6 minutes, $p < 0.05$). The particle growth rate slowed down in presence of HPMC. However, the growth rate was not decreased further as the HPMC concentration increased (0.01-1 %w/v). A clear difference was observed in particle size when comparing tadalafil particles with and without PI (*Fig. 1 and 2*).

CONCLUSION

Tadalafil, in presence of HPMC, shows a prolonged induction time and a reduced particle growth rate. Increasing HPMC concentrations resulted in increased induction time, but not in changes in the crystal growth rate. This method is a promising tool for comparing the effectiveness of PI on poorly soluble drugs.

ACKNOWLEDGEMENTS

This study was conducted as part of the Oral Biopharmaceutics Tools (ORBITO) project (<http://www.orbitoproject.eu>), funded by the Innovative Medicines Initiative (IMI) Joint Undertaking under Grant Agreement No. 115369.